SUNTEXT REVIEWS

SunText Review of Pediatric Care
ISSN: 2766-5216

Open Access
Research Article
Volume 4:1

Cerebrovascular Accidents in Sickle Cell Disease; Risk Factor and Prevention: Systemic Review

Elfaki AAA[1,*] and Ahmed HAA[2]

[1]*Faculty of Medicine, International University of Africa Sudanese Medical Research Association (SMRA), Sudan*
[2]*Faculty of Medicine, Sudan International University Sudanese Medical Research Association (SMRA), Sudan*

[*]**Corresponding author:** Elfaki AAA, Faculty of Medicine, International University of Africa Sudanese Medical Research Association (SMRA), Sudan

Received date: 27 October 2023; **Accepted date:** 29 November 2023; **Published date:** 05 December 2023

Citation: Elfaki AAA, Ahmed HAA (2023). Cerebrovascular Accidents in Sickle Cell Disease; Risk Factor and Prevention: Systemic Review. SunText Rev Pediatr Care 4(1): 142.

DOI: https://doi.org/10.51737/2766-5216.2023.042

Copyright: © 2023 Elfaki AAA, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Abstract

Sickle cell disorder one of well-known problems affecting children. Cerebrovascular accidents (CVA) is the most complication and fundamental presenting factor of sickle cell anaemia. There are many hazard factors contributing in development of Sickle cell anaemia as nicely as lead to strokes as complications. This assessment aiming to talk about the preventive methods of cerebrovascular accidents (CVA) in affected person with Sickle Cell Disease (SCD), and to evaluate the most common danger elements of the disease. The evaluation was conducted with the aid of searching in Medline, Embassy, Web of Science, Science Direct, BMJ journal, and Google Scholar for, researches, evaluate articles, and reports, posted over the previous years. Were searched up to June 2021 for published and unpublished studies and without language restrictions, we randomly selected one or two to keep away from repetitive outcomes in similar findings items. Based on the findings we located that SS phenotype and growing older in SCD sufferers are the most vital risk and primary leading cause of stroke. The most appropriate preventive strategies of cerebrovascular accident in patient with sickle mobile disorder is blood transfusion, also is located that hydroxyurea has a role in lowering chance of Stroke, and it is want to be applied specially to stroke prevention in randomized trials.

Keywords: Cerebrovascular accidents; Sickle cell disease; Risk factor; Prevention

Introduction

One defective structure of the gene, humans with the sickle cell trait make each normal haemoglobin and sickle cell haemoglobin. Their blood may contain some sickle cells; however, they generally don't have symptoms. They're carriers of the disease, however, which capability they can bypass the gene to their children [2]. Most frequent Types of SCD Include: a. HbSS: People who have this form of SCD inherit two sickle cell genes (“S”), one from each parent. This is usually referred to as sickle cell anaemia and is commonly the most severe structure of the disease. b. HbSC: People who have this structure of SCD inherit a sickle cell gene (“S”) from one parent and from the other parent a gene for an abnormal haemoglobin called “C”. Haemoglobin is a protein that allows red blood cells to carry oxygen to all components of the body. This is typically a milder form of SCD. c. HbS beta thalassemia: People who have this form of SCD inherit one sickle cell gene (“S”) from one parent and one gene for beta thalassemia, another type of anaemia, from the different parent. There are two types of beta thalassemia: “0” and “+”. Those with HbS beta 0- thalassemia typically have an extreme form of SCD. People with HbS beta +-thalassemia tend to have a milder form of SCD [3].

Sickle Cell Trait (SCT): (HbAS): People who have SCT inherit one sickle cell gene (“S”) from one parent and one normal gene (“A”) from the other parent. This is known as sickle cell trait (SCT). People with SCT typically do no longer have any of the signs of the disease and live a normal life, but they can pass the trait on to their children. Additionally, there are a few, uncommon health problems that may also probably be related to sickle cell. There's no cure for most people with sickle cell anaemia. But treatments can relieve pain and help prevent complications

associated with the disease [3]. Most frequent complication of Sickle Cell Anaemia is cerebrovascular accidents (CVA) in Children. Children 5%: Highest risk at ages two to 5 years old with Sickle Cell Anaemia. Clinically obvious stroke occurs in 10% by way of age 5 years old [4]. Silent cerebral infarct happens in up to 25% of youth through age 6 years old and it can also end result in gradual cognitive decline. Young adult (age 20): 11%. Age forty-five years old: 24%. Lifetime risk: 25 % [5]. The highest charges of prevalence of CVA (4.01%) and incidence (0.61 per 100 patient-years) have been in sickle cell anaemia (SS) patients, however CVA took place in all frequent genotypes. The incidence of infective CVA used to be lowest in SS sufferers 20 to 29 years of age and higher in youth and older sufferers [6]. This study aims to determine the preventive techniques of cerebrovascular accidents in patients with Sickle cell anaemia, in addition to consider the risk factors the lead to develop cerebrovascular accidents in Sickle cell patients.

Materials and Methods

This an overview was conducted in July 2021 below the preferred reporting objects for systematic reviews and meta-analyses (PRISMA) declaration standards. We reviewed all the topics on the prevention and risk factors of cerebrovascular accidents on Sickle Cell patients.

***Table 1*: *Selected data*.**

Author/s and Year	Design	Sample	Finding	Level of evidence
Ohene-Fremponget al. 1998	Longitudinal clinical Study Oct 1978 – Sept 1988	4082 patients	Highest rate Prevalence of CVA in SCD is 4.01%, SS phenotype patients has the highest risk	Level 1 evidence
Adams RJ, McKieVC, Hsu L, et al. 2001	Section Editor: IRA SHOULSON,MD		Two newer treatments for SCD, hydroxyurea therapy and BMT, need to be applied specifically to stroke preventionin randomized trials.	Level 3 evidence
Adetola A. Kessler. 2016	Section Editor: Craig M. Kessler,MD			Level 3 evidence
Allison A King etal. 2014	Multicenter, cross-sectionalstudy	150 children	FSIQ in children with sickle cell anemia is beat accounted for by a multivariate model that includes both biologic and Socioenvironmental factors.	Level 2 evidence

Our search was once done barring any language restrictions. Then information used to be extracted on study design, and key outcome on prevention and risk factors of cerebrovascular accidents on Sickle Cell patients. The selected studies have been summarized and unreproducible studies were excluded (Table 1). Studies have been rated as being high great with the aid of an established contrast process primarily based on the Dynamic criteria.

Inclusion criteria

Current prevention strategies of Cerebrovascular accidents on patients suffering from SCD, and the most frequent risk factors leading to advance Cerebrovascular accidents in sickle cell patients.

Exclusion criteria

Irrelevant articles not related to the purpose of this review and articles that did not meet the inclusion criteria in this review.

Data extraction and analysis

Information concerning to every of the systematic review query elements used to be extracted from the studies and collated in qualitative tables. Then direct evaluation of the research associated to cerebrovascular accidents and Sickle Cell Disease on prevention and risk factors, had been Done.

Results and Discussion

Strokes take place in approximately about 5% to 10% of youngsters with sickle cell disease (SCD). Patients with Genotype SS or Sβ0 thalassemia of SCA are the most frequent current with strokes. Ischemic stroke has a bimodal distribution, being more common in children and older adults and less common in adults aged 20 to 29 years, while hemorrhagic stroke has been proven to be most familiar in the 20- to 29-year age group [7]. The most many times diagnosed cause of neurologic damage is Silent cerebral infarcts (ischemic lesions that are detected with magnetic resonance imaging (MRI)). Silent strokes are cerebral infarcts that have a sign abnormality measured at a minimal of three mm (visible on fluid-attenuated inversion restoration MRI in both axial and coronal views), do not cause abnormalities that are revealed on neurologic examination, and are associated with cognitive difficulties [8]. Among adults with SCA, the mentioned risk for overt stroke is 11% through 20 years of age and 24% by forty-five years of age, [7] while for younger patients affecting by using SCA, the cumulative risk for stroke is 11.5% by 18 years of age and 12.8% by way of 20 years of age [9]. Silent cerebral infarcts are most frequent in SCA which occur in 20% to 40% of teens with SCD. SS phenotype in patients of SCA are at the highest chance for stroke, for each overt and silent strokes, and this threat continues which growing with age. While age increase danger of stroke in different phenotypes, which can be problematic via other known risk factors for stroke, such as hypertension, renal disease, diabetes mellitus, atrial fibrillation, and hyperlipidaemia [10]. Chronic blood transfusion therapy has been proven to decrease the annual risk for stroke from 10% to less than 1%, which prevent stroke via 92% [11]. From 1970s to 1980s, clinical series from countless facilities indicated that children with SCD and stroke had a very excessive early (3 years) recurrent stroke risk [12]. And that if they receive transfusion remedy this threat reduced [13]. In most cases, the transfusion programs have been sufficient to reduce total sickle cell haemoglobin values to less than 30% of the whole haemoglobin values. According to STOP (Stroke Prevention Trial in Sickle Cell Anemia), the transcranial Doppler ultrasonography (TCD) velocities of many patients undergoing transfusion reverted from excessive hazard to curiously low risk (170 cm/s, approximately 53%) or intermediate risk (170-199 cm/s, about 17%) [14]. However, TWiTCH (Transcranial Doppler With Transfusions Changing to Hydroxyurea), a study carried out by way of Ware and colleagues, showed that in high threat children with SCA and abnormal TCD velocities who have obtained transfusions for at least 1 year and have no MRA-defined extreme vasculopathy, hydroxycarbamide treatment can be substituted for chronic transfusions to maintain TCD velocities and reduce the risk for predominant stroke [15]. Hydroxyurea was once only chemotherapeutic agent which accredited for the treatment of SCD. The double-blind, placebo-controlled find out about of hydroxyurea therapy in 299 adults with SCD determined that there is reduction in painful episodes, acute chest syndrome, need for hospitalization, and blood transfusions grew to become evident [16]. No study has addressed the problem of whether or not hydroxyurea therapy has efficacy in stroke prevention in a controlled fashion. Ware et al [17] suggested the results of secondary stroke treatment with hydroxyurea and phlebotomy in 16 younger patients in whom transfusion was no longer an option. Their results of a 19% recurrent tournament incidence are encouraging however want to be in contrast with an excellent control. In this single report, the sample size used to be small and there were no controls or and randomization.

Conclusion

The study come out that SS phenotype in patients of SCA are the most essential risk and primary leading cause of stroke. Increasing in age of SCD patients can be considered as the second necessary risk factor for developing Stroke. One of the important complications of SCD are Strokes, and it is indispensable and affect teens life; While the most fabulous preventive strategies of cerebrovascular accident in affected person with sickle mobile disease is blood transfusion. Hydroxyurea therapy found to have an impact in decreasing mortality rate from strokes. Two more modern treatments for SCD, hydroxyurea therapy and BMT, want to be applied specifically to stroke prevention in randomized trials.

Conflict of Interest

The authors of this article hasn’t receive any support of this work and it was completely self-funded.

References

1. Field JJ, Vichinsky EP, DeBaun MR. Overview of the management and prognosis of sickle cell disease. 2019.
2. National Heart L, and Blood Institute. Sickle cell disease. 2019.
3. Center of Disease prevention and control CDC Web page: National Center on Birth Defects and Developmental Disabilities; 2020.

4. Baskin MN, Goh XL, Heeney MM, Harper MB. Bacteremia risk and outpatient management of febrile patients with sickle cell disease. Pediatrics. 2013; 131: 1035-1041.
5. Cerebrovascular Accident in Sickle Cell Anemia Web page: Hematology and Oncology Book, Hemoglobinopathies. 2018.
6. Ohene-Frempong K , Sleeper LA, Miller ST, Embury S, Moohr JW, Wethers DL, et al. Cerebrovascular accidents in sickle cell disease: rates and risk factors. Blood. 1998; 91: 288-294.
7. King AA, Rodeghier MJ, Compas BE, Strouse JJ, Casella JF, Noetzel MJ, et al. Parent education and biologic factors influence on cognition in sickle cell anemia. Am J Hematol. 2014; 89: 162-167.
8. Telfer P, Coen P, Chakravarty S, Wilkey O, Evans J, Newell H, et al. . Clinical outcomes in children with sickle cell disease living in England: a neonatal cohort in East London. Haematologica. 2007; 92: 905-912. .
9. Kassim AA. Prevention and tretment of Stroke in patients with sikle Cell Diseases. Clinical Advance in hematology and oncology. 2016; 14.
10. Adams RJ, McKie VC, Hsu L, Files B, Vichinsky E, Pegelow, et al. Prevention of a first stroke by transfusions in children with sickle cell anemia and abnormal results on transcranial Doppler. Ultrasonography. N Engl J Medicine. 1998; 339: 5-11.
11. Powears D, Wilson B, Lmbus C, Pegelow C, Allen J. The natural history of stroke in sickle cell disease. Am J Med. 1978; 65: 461-471.
12. Russell MO. Goldberg HL, Hodson A, Kim HC, Halus J, Reivich M, et al. Effect of transfusion therapy on arteriography abnormalities and on recurrence of stroke in sickle cell disease. Blood. 1984; 63: 162-169.
13. Lee MT, Piomelli S, Granger S, Miller ST, Harkness S, Brambilla DJ, et al. STOP Study Investigators. Stroke Prevention Trial in Sickle Cell Anemia (STOP): extended follow-up and final results. Blood. 2006; 108: 847-852.
14. Ware RE, Davis BR, Schultz WH, Brown RC,Aygun B,Samaik S, et al. Hydroxycarbamide versus chronic transfusion for maintenance of transcranial doppler flow velocities in children with sickle cell anaemia- TCD With Transfusions Changing to Hydroxyurea (TWiTCH): a multicentre, open-label, phase 3, non- inferiority trial. Lancet. 2016; 387: 661-670.
15. Charache S, Terrin ML, Moore RD, Dover Gj. Barton FB. Effect of hydroxyurea on the frequency of painful crisis in sickle cell anemia. N Engl Medicine. 1995; 332: 1317-1322.
16. Ware RE, Zimmerman SA, Schultz WH. Hydroxyurea as an alternative to blood transfusions for the prevention of recurrent stroke in children with sickle cell disease. Blood.1999; 94: 3022-3026.