



# A Rare Vascular Complication in a Covid-19 patient: A Case Report of Free-Floating Carotid Thrombus

Fatma BS\*, Chaker J, Khedija S, Mouna B, Nadia A and Amine J

Director, Abderrahman Mami Hospital Ariana, Tunisia

\*Corresponding author: Fatma BS, Abderrahman Mami Hospital Ariana Tunisia; E-mail: [fatmabensaida@gmail.com](mailto:fatmabensaida@gmail.com)

## Abstract

It is now admitted that Coronavirus is related to vascular complications especially thrombotic ones. Mechanism of thrombotic complications is not fully understood. Free-floating carotid thrombus is a rare condition and it has been found in numerous cases of Covid-19 patients. Data related to management of this floating clot is limited. Anticoagulation remains an important pillar of treatment. We report in this work, a case of a 59-year-old male, hospitalized twice for management of Covid-19 infection in January and March 2021. A free-floating carotid thrombus was discovered and a medical treatment was initiated.

**Keywords:** COVID-19; Free-floating thrombus; Anticoagulation

## Introduction

Since its emergence in March 2020, Coronavirus symptoms and complications are constantly being discovered especially the vascular ones. It can be explained by the inflammation process, the immobilization and excessive activation of coagulation system. These factors lead to thrombi formation in either arterial and venous systems. The free-floating thrombus in carotid artery is a described complication in covid-19 patients. We report the case of a patient hospitalized for the management of a COVID-19 pneumonia with a symptomatic free-floating carotid thrombus.

## Case Description

We are reporting the case of a 59-year-old male patient, with cardiovascular risk factors: Weaned tobacco user for four years, Hypertension and Diabetes on insulin. He contracted Covid-19 in January 2021 and was hospitalized for Oxygen therapy. He was put on Azithromycin and Imipenem for ten days associated to prophylactic anticoagulation therapy. During hospitalization, he presented a transient ischemic attack (TIA): A paresis of the left upper limb. Brain imaging was normal. It was decided to put the patient on therapeutic dose of anticoagulation. Few days later, the patient was discharged. On March 2021, he was hospitalized for a

covid-19 reinfection with need to oxygen therapy. One week after being hospitalized, he presented a stroke with left hemiplegia and aphasia. CT scan confirmed the ischemic nature of the stroke. CT angiography showed a floating thrombus in the right internal carotid artery (Figure 1).



**Figure 1:** CT ANGIOGRAPHY showing the free-floating thrombus in the right internal carotid artery.

Received date: 10 February 2026; Accepted date: 02 March 2026; Published date: 10 March 2026

**Citation:** Fatma BS, Chaker J, Khedija S, Mouna B, Nadia A, Amine J (2026) A Rare Vascular Complication in a Covid-19 patient: A Case Report of Free-Floating Carotid Thrombus. SunText Rev Case Rep Image 7(1): 174.

**DOI:** <https://doi.org/10.51737/2766-4589.2026.174>

**Copyright:** © 2026 Fatma BS, et al. This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.

Our decision was to maintain the therapeutic dose of anticoagulation for two weeks and then reassess for a possible surgical management. Two weeks after complete anticoagulation, CT angiography was done showing a complete dissolution of the floating clot. The patient was discharged after few days.

## Discussion

Carotid free-floating thrombus (CFFT) is uncommon [1]. This entity was discovered and described in 1966 by Ehrenfeld. Its occurrence in 0,4% of cases in non-Covid-19 patients [2]. Since the emergence of the COVID-19 infection, we are describing the case N-16 of a free-floating clot in the carotid artery complicating SARS-Cov 2 infection. In our knowledge, this is the first described case in TUNISIA. CFFT is defined as a blood clot attached to the arterial wall with blood flow in distality [3]. The main etiology of this state, is migration or rupture of an atherosclerotic plaque. Other etiologies are rare including arterial dissection, aneurysm, prothrombotic factors such as inflammatory and infectious diseases [4]. The internal carotid artery is commonly the most affected with an estimated frequency of 7% [5]. It's now known that Covid-19 infection is related to an hyperinflammation state. Indeed, higher levels of WBC count and neutrophil-to-lymphocyte ratio with important rates of CRP and CPK levels were found in Sars cov-2 patients compared to other patients. This hyperinflammation can be explained by an abnormal circulating immune complex formation in response to COVID-19 infection [6]. This Inflammation state associated with the platelets activation due to infection are probably involved in the pathogenesis of hypercoagulability leading to thrombi formation. Furthermore, disseminated intravascular coagulation is more frequent in Covid-19 patients with a rate of 8,7% [7]. Other hypothesis suggested that there is an endothelial dysfunction caused by invasion of endothelial cells by the virus [8]. This hypercoagulability leads to a higher prothrombotic state in Sars-cov 2 patients explaining the increasing frequency of vessel occlusions even without atherosclerotic disease.

In literature, the COVID-19 patients with artery thrombosis characteristics are male gender, obesity and less cardiovascular factors compared to other patients [9]. In most cases, CFFT is symptomatic with a history of ischemic stroke [10]. The rate of stroke in Covid-19 patients is about 1% [11]. The feature of this type of thrombus lies in its extreme fragility leading to recurrent cerebral embolism. CT angiography is the gold standard in the diagnosis of CFFT finding a specific sign called the donut sign [12]. The treatment is either a surgical management or optimal medical strategy. Hosseini et al. reported a complete resolution of the clot three months after the initiation of the anticoagulation therapy [13]. In a study published in 2013, 24 patients were included with the diagnosis of intraluminal carotid thrombus [14]. All of them had an anticoagulation therapy first, ten of them had a

delayed revascularization. The results reported of all patients with the anticoagulation therapy in primary intention were excellent with no ischemic nor hemorrhagic stroke. When endarterectomy is indicated, the surgery should be done a maximum of two weeks after the patients last symptoms [15]. Endovascular approach can also be considered in CFFT with stenting, aspiration or reversal of flow [10]. The major problem of arterial thrombosis in Covid-19 patients remains the risk of re-thrombosis despite anticoagulation. In fact, cases of recurrence have been reported in literature and even in our daily practice, patients who underwent surgical management for artery thrombosis and then have had an anticoagulation, presented a re-thrombosis.

## Conclusion

COVID-19 is associated with a higher risk of thrombotic events. Free-floating carotid thrombus is uncommon but have been reported in now 16 cases in COVID-19 patients since the virus emergence in March 2020. The management strategy of this state remains limited and is based on case reports. Anticoagulation in primary intention seems to have good results. A re-thrombosis should be kept in mind even after full treatment and a prophylactic strategy must be considered.

## References

1. Lane TRA, Shalhoub J, Perera R, Mehta A, Ellis MR, Sandison A, et al. Diagnosis and surgical management of free-floating thrombus within the carotid artery. *Vasc Endovascular Surg.* 2010; 44: 586-593.
2. Choi H-Y, Ye BS, Ahn SH, Cho HJ, Kim D-J, Kim SM, et al. Characteristics and the fate of intraluminal thrombus of the intracranial and extracranial cerebral arteries in acute ischemic stroke patients. *Eur Neurol.* 2009; 62: 72-78.
3. Bhatti AF, Leon LR, Labropoulos N, Rubinas TL, Rodriguez H, Kalman PG, et al. Free-floating thrombus of the carotid artery: Literature review and case reports. *J Vasc Surg. janv* 2007; 45: 199-205.
4. Fridman S, Lownie SP, Mandzia J. Diagnosis and management of carotid free-floating thrombus: A systematic literature review. *Int J Stroke.* 2019; 14: 247-256.
5. Singh RJ, Chakraborty D, Dey S, Ganesh A, Al Sultan AS, Eesa M, et al. Intraluminal thrombi in the cervico-cephalic arteries: clinical-imaging manifestations, treatment strategies, and outcome. *stroke.* 2019; 50: 357-364.
6. Indes JE, Koleilat I, Hatch AN, Choinski K, Jones DB, Aldailami H, et al. Early experience with arterial thromboembolic complications in patients with COVID-19. *J Vasc Surg.* 2021; 73: 381-389.
7. Yaghi S, Ishida K, Torres J, Mac Grory B, Raz E, Humbert K, et al. SARS-CoV-2 and stroke in a New York healthcare system. *Stroke.* 2020; 51: 2002-2011.
8. Gomez-Arbelaez D, Ibarra-Sanchez G, Garcia-Gutierrez A, Comanges-Yeboles A, Ansuategui-Vicente M, Gonzalez-Fajardo JA.



## SUNTEXT REVIEWS

- COVID-19-related aortic thrombosis: a report of four cases. *Ann Vasc Surg.* 2020; 67: 10-13.
9. Cancer-Perez S, Alfayate-Garcia J, Vicente-Jimenez S, Ruiz-Munoz M, Dhimes-Tejada FP, Gutierrez-Baz M, et al. Symptomatic common carotid free-floating thrombus in a covid-19 patient, case report and literature review. *Ann Vasc Surg.* 2021; 73: 122-128.
  10. Tolaymat B, Irizarry K, Reif M, Drucker CB, Aicher BO, Sarkar R, et al. Considerations beyond stenosis for carotid endarterectomy in treating free-floating thrombus of the carotid artery. *Ann Vasc Surg.* 2019; 60: 221-228.
  11. Mowla A, Sizdahkhani S, Sharifian-Dorche M, Selvan P, Emanuel BA, Tenser MS, et al. Unusual pattern of arterial macrothrombosis causing stroke in a young adult recovered from COVID-19. *J Stroke Cerebrovasc Dis.* 2020; 29: 105353.
  12. for the Calgary CTA Study Group, Menon BK, Singh J, Al-Khataami A, Demchuk AM, Goyal M. The donut sign on CT angiography: an indicator of reversible intraluminal carotid thrombus? *Neuroradiol.* 2010; 52: 1055-1056.
  13. Hosseini M, Sahajwani S, Zhang J, Toursavadkahi S, Ucuzian AA. Delayed stroke after hospitalization for coronavirus disease 2019 pneumonia from common and internal carotid artery thrombosis. *J Vasc Surg Cases Innov Tech.* 2021; 7: 40-45.
  14. Vellimana AK, Kadkhodayan Y, Rich KM, Cross DT, Moran CJ, Zazulia AR, et al. Symptomatic patients with intraluminal carotid artery thrombus: outcome with a strategy of initial anticoagulation: Clinical article. *J Neurosurg.* 2013; 118: 34-41.
  15. Sallustio F, Di Legge S, Marziali S, Ippoliti A, Stanzione P. Floating carotid thrombus treated by intravenous heparin and endarterectomy. *J Vasc Surg.* 2011; 53: 489-491.